




Διαχείριση του Διαβητικού Ασθενούς με Σήψη



Ερυφίλη Χατζηαγγελάκη
**Καθηγήτρια Παθολογίας-
Μεταβολικών Νοσημάτων**
Β' Προπ. Παθολογική Κλινική,
Μονάδα Έρευνας & Διαβητολογικό Κέντρο
Πανεπιστημιακό Γ.Ν "Αττικόν"
Υπεύθυνη Διαβητολογικού Κέντρου

***Visiting Professor, German Diabetes Center
(DDZ) at Heinrich Heine University of
Düsseldorf, Germany***
***Vice President, Central European Diabetes
Association (CEDA)***

Prevalence of diabetes (2017 and 2045)



The age group 65–79 years shows the highest diabetes prevalence in both women and men
International Diabetes Federation. IDF Diabetes Atlas. 8th edn. Brussels, Belgium: International Diabetes Federation, 2017.
Available at: <http://www.diabetesatlas.org> Accessed November 2018

DM: 7th leading cause of death

CV
Death

1.7 x more likely

MI
Hospital-
ization

1.8 x more likely

Stroke
Hospital-
ization


1.5 x more likely

CKD

44% of all new cases

Every 6 seconds, 1 person dies from diabetes-related complications

Diabetes significantly increases the risk of...



**Heart
disease by
2-4 fold**



**Stroke
by > 2-4 fold**



...there will be 1104 new cases of diabetic retinopathy, which can lead to vision loss




...133 patients will start dialysis



...180 patients will have an amputation


Various parameters impact T2D pathophysiology

THE 'OMINOUS OCTET'



Drugs with different mechanisms of action are required to address the numerous Type 2 diabetes pathophysiological defects

The ominous octet




HGP, hepatic glucose production


1. DeFronzo RA. *Diabetes*. 2009; 58:773–795. 2. Sharma MD *et al.* *Diabetes Obes Metab*. 2015; 17:616–621.

3. Abdul-Ghani M *et al.* *Diabetes Care*. 2015; 38:373–375.


Η Πρωιμότερη και Ενδεδειγμένη Παρέμβαση Μπορεί να Βελτιώσει τις Πιθανότητες των Ασθενών να Επιτύχουν το Στόχο



Unmet Needs in Diabetes Care



Multifactorial approach improves outcomes



- The ADA recommends that most adults with diabetes achieve the following targets:¹
 - HbA_{1c}: <7.0% (53 mmol/mol)
 - Physical activity: ≥150 min per week*
 - Blood pressure: <140/90 mmHg
 - Triglycerides: <150 mg/dL (1.7 mmol/L)
 - HDL-C: ≥40–50 mg/dL (1–1.3 mmol/L)[†]
 - LDL-C: <100 mg/dL (2.6 mmol/L)

Primary prevention methods only - platelet inhibition recommended as secondary prevention.

*Physical activity of moderate to vigorous intensity, spread over at least 3 days/week. [†]ADA guidelines suggest 40 mg/dL for men, 50 mg/dL for women.

ACE-I, angiotensin-converting enzyme inhibitor; ADA, American Diabetes Association; ARB, angiotensin receptor blocker; BP, blood pressure; HDL-C, high-density lipoprotein cholesterol; LDL-C, low-density lipoprotein cholesterol. 1. American Diabetes Association. *Diabetes Care* 2019;42(Suppl. 1):S1–S3.



Management of Hyperglycemia in Type 2 Diabetes, 2018. A Consensus Report by the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD)

https://doi.org/10.2337/181122

Diabetologia
https://doi.org/10.1007/s00125-018-4729-5

CONSENSUS REPORT



Management of hyperglycaemia in type 2 diabetes, 2018. A consensus report by the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD)

Melanie J. Davies^{1,2} · David A. D'Alessio³ · Judith Fradkin⁴ · Walter N. Kernan⁵ · Chantal Mathieu⁶ · Gertrude Mingrone^{7,8} · Peter Rossing^{9,10} · Apostolos Tzazpas¹¹ · Deborah J. Wexler^{12,13} · John B. Buse¹⁴

© European Association for the Study of Diabetes and American Diabetes Association 2018

Abstract

The American Diabetes Association and the European Association for the Study of Diabetes convened a panel to update the prior position statements, published in 2012 and 2015, on the management of type 2 diabetes in adults. A systematic evaluation of the literature since 2014 informed new recommendations. These include additional focus on lifestyle management and diabetes self-management education and support. For those with obesity, efforts targeting weight loss, including lifestyle, medication and surgical interventions, are recommended. With regards to medication management, for patients with clinical cardiovascular disease, a sodium-glucose cotransporter-2 (SGLT2) inhibitor or a glucagon-like peptide-1 (GLP-1) receptor agonist with proven cardiovascular benefit is recommended. For patients with chronic kidney disease or clinical heart failure and atherosclerotic cardiovascular disease, an SGLT2 inhibitor with proven benefit is recommended. GLP-1 receptor agonists are generally recommended as the first injectable medication.

Keywords Cardiovascular disease · Chronic kidney disease · Costs · Glucose-lowering therapy · Guidelines · Heart failure · Hyperglycaemia · Patient-centred care · Type 2 diabetes mellitus · Weight management

Abbreviations

| | | | |
|--------|---|------------------|--|
| ARR | Absolute risk reduction | DKA | Diabetic ketoacidosis |
| ASCVD | Atherosclerotic cardiovascular disease | DPP-4 | Dipeptidyl peptidase-4 |
| CANVAS | Canagliflozin Cardiovascular Assessment Study | DPP-4i | Dipeptidyl peptidase-4 inhibitor |
| CKD | Chronic kidney disease | DSMES | Diabetes self-management education and support |
| CVD | Cardiovascular disease | EMPA-REG OUTCOME | Empagliflozin, Cardiovascular Outcome Event Trial in Type 2 Diabetes Mellitus Patients |
| CVOT | Cardiovascular outcomes trial | ESRD | End-stage renal disease |
| | | EXSCEL | Exenatide Study of Cardiovascular Event Lowering |

M. J. Davies and J. B. Buse were co-chairs for the Consensus Statement Writing Group. D. A. D'Alessio, J. Fradkin, W. N. Kernan and D. J. Wexler were the writing group members for the ADA. C. Mathieu, G. Mingrone, P. Rossing and A. Tzazpas were writing group members for the EASD.

This article is being simultaneously published in *Diabetes Care* and *Diabetologia* by the American Diabetes Association and the European Association for the Study of Diabetes.

✉ Melanie J. Davies
Melanie.davies@ahd-trnab.ch

Extended author information available on the last page of the article

Melanie J. Davies^{1,2} · David A. D'Alessio³ · Judith Fradkin⁴ · Walter N. Kernan⁵ · Chantal Mathieu⁶ · Gertrude Mingrone^{7,8} · Peter Rossing^{9,10} · Apostolos Tzazpas¹¹ · Deborah J. Wexler^{12,13} · John B. Buse¹⁴

for the Study of Diabetes, published in *Diabetologia*. These self-management weight loss, amended. With cardiovascular a a glucagon-like benefit is recom heart failure and proven benefit is noted as the first

of glycemia in and maintaining management and is summarized in recommendations and secondary diabetes,

of this position conducted an rd meta-analysis,

t, published online October 4, 2018

¹Diabetes Research Centre, University of Leicester, Leicester, U.K.

²Leicester Diabetes Centre, Leicester General Hospital, Leicester, U.K.

³Department of Medicine, Duke University School of Medicine, Durham, NC

⁴National Institute of Diabetes and Digestive and Kidney Disorders, National Institutes of Health, Bethesda, MD

⁵Department of Medicine, Yale School of Medicine, New Haven, CT

⁶Clinical and Experimental Endocrinology, UZ Ghent/Leuven, KU Leuven, Leuven, Belgium

⁷Department of Internal Medicine, Catholic University, Rome, Italy

⁸Diabetes and Nutritional Sciences, King's College London, London, U.K.

⁹Steno Diabetes Center Copenhagen, Gentofte, Denmark

¹⁰University of Copenhagen, Copenhagen, Denmark

¹¹Second Medical Department, Aristotle University Thessaloniki, Thessaloniki, Greece

¹²Department of Medicine and Diabetes Unit, Massachusetts General Hospital, Boston, MA

¹³Harvard Medical School, Boston, MA

¹⁴Department of Medicine, University of North Carolina School of Medicine, Chapel Hill, NC

Corresponding author: John B. Buse, jbuse@med.unc.edu.

M.J.D. and J.B.B. were co-chairs for the Consensus Statement Writing Group. D.A.D.A., J.F., W.N.K., and D.J.W. were the writing group members for the American Diabetes Association. C.M., G.M., P.A., and A.T. were writing group members for the European Association for the Study of Diabetes. This article is being simultaneously published in *Diabetes Care* and *Diabetologia* by the American Diabetes Association and the European Association for the Study of Diabetes.


© 2018 American Diabetes Association and European Association for the Study of Diabetes. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. More information is available at <http://www.diabetologyandendocrinology.com>.

ADA/EASD 2018 Consensus Report

ADA/EASD 2018 consensus for glucose-lowering medication in T2D

**FIRST-LINE THERAPY IS METFORMIN AND COMPREHENSIVE LIFESTYLE (INCLUDING WEIGHT MANAGEMENT AND PHYSICAL ACTIVITY)
IF HbA_{1c} ABOVE TARGET PROCEED AS BELOW**


To avoid clinical inertia reassess and modify treatment regularly (3-6 months)




*Proven CVD benefit means it has label indication of reducing CVD events. For GLP-1RA strongest evidence for liraglutide>semaglutide>exenatide extended release. For SGLT-2i evidence mostly stronger for empagliflozin>canagliflozin; †Be aware that SGLT-2i vary by region and individual agent with regard to indicated level of eGFR for initiation and continued use; ‡Both empagliflozin and canagliflozin have shown reduction in HF and reduction in CKD progression in CVOTs; §Degludec or U100 glargine have demonstrated CVD safety; ¶Low dose may be better tolerated though less well studied for CVD effects; ||Choose later generation SU with lower risk of hypoglycaemia; **Semaglutide>liraglutide>dulaglutide>exenatide>lixisenatide; ††If no specific comorbidities (i.e. no established CVD, low risk of hypoglycaemia and lower priority to avoid weight gain or no weight-related comorbidities); ***Consider country- and region-specific cost of drugs. In some countries, TZDs relatively more expensive and DPP-4i relatively cheaper. Davies MJ et al. *Diabetologia* 2018;61:2461-2498; Davies MJ et al. *Diabetes Care* 2018;41:2669-2701

WITHOUT ESTABLISHED ASCVD OR CKD


COMPELLING NEED TO MINIMISE HYPOGLYCAEMIA




COMPELLING NEED TO MINIMISE WEIGHT GAIN OR PROMOTE WEIGHT LOSS



COST IS A MAJOR ISSUE⁹⁻¹⁰



INTENSIFYING TO INJECTABLE THERAPIES





Consensus recommendation

- In patients who need the greater glucose-lowering effect of an injectable medication, GLP-1 receptor agonists are the preferred choice to insulin. For patients with extreme and symptomatic hyperglycemia, insulin is recommended

INITIATION FOR BASAL
• Start 10 IU a day OR 0.1–0.2 IU/kg a day

ADA-EASD 2018 consensus


In patients with established ASCVD



* Proven CVD benefit means it has label indication of reducing CVD events. For GLP-1RA strongest evidence for liraglutide>semaglutide>exenatide extended release. For SGLT-2i evidence modestly stronger for empagliflozin>canagliflozin; [†]Be aware that SGLT-2i vary by region and individual agent with regard to indicated level of eGFR for initiation and continued use; [‡]Both empagliflozin and canagliflozin have shown reduction in HF and reduction in CKD progression in CVOTs; [§]Degludecor U 100 glargine have demonstrated CVD safety; [¶]Low dose may be better tolerated though less well studied for CVD effects; ^{||}Choose later generation SU with lower risk of hypoglycaemia

ADA-EASD 2018 consensus

In patients with established HF or CKD



- **SGLT-2is preferred** over GLP-1RAs as significant, consistent reductions in **hospitalisation for HF and CKD** progression have been seen in SGLT-2i trials

Evidence Hierarchy



Evidence Hierarchy



*Proven CVD benefit means that it has label indication of reducing CVD events. For GLP-1RAs, strongest evidence for liraglutide > semaglutide > exenatide extended release. For SGLT-2is, evidence modestly stronger for empagliflozin > canagliflozin; [‡]Be aware that SGLT-2is vary by region and individual agent with regard to indicated level of eGFR for initiation and continued use; [‡]Both empagliflozin and canagliflozin have shown reduction in HF and reduction in CKD progression in CVOTs; [§]Degludec or U100 glargine have demonstrated CVD safety; ^{||}Choose later-generation SU with lower risk of hypoglycaemia; [#]Caution with GLP-1RA in ESRD

2019 ADA Standards of Care

Table 9.1—Drug-specific and patient factors to consider when selecting antihyperglycemic treatment in adults with type 2 diabetes

| | Efficacy | Hypoglycemia | Weight change | CV effects | | Cost | Oral/SQ | Renal effects | | Additional considerations |
|--------------------------|--------------|--------------|-------------------------------------|--|--|------|---------|--|---|--|
| | | | | ASCVD | CHF | | | Progression of DKD | Dosing/use considerations* | |
| Metformin | High | No | Neutral (potential for modest loss) | Potential benefit | Neutral | Low | Oral | Neutral | <ul style="list-style-type: none"> Contraindicated with eGFR <30 | <ul style="list-style-type: none"> Gastrointestinal side effects common (diarrhea, nausea) Potential for B12 deficiency |
| SGLT-2 inhibitors | Intermediate | No | Loss | Benefit: empagliflozin, canagliflozin | Benefit: empagliflozin, canagliflozin  | High | Oral | Benefit: canagliflozin, empagliflozin  | <ul style="list-style-type: none"> Renal dose adjustment required (canagliflozin, dapagliflozin, empagliflozin, ertugliflozin) | <ul style="list-style-type: none"> FDA Black Box: Risk of amputation (canagliflozin) Risk of bone fractures (canagliflozin) DKA risk (all agents, rare in T2DM) Genitourinary infections Risk of volume depletion |
| GLP-1 RAs | | | | Benefit: liraglutide† > semaglutide > exenatide extended release | | | | | <ul style="list-style-type: none"> Caution when initiating or increasing dose due to potential risk of acute kidney injury | <ul style="list-style-type: none"> Gastrointestinal side effects common (nausea, vomiting, diarrhea) Injection site reactions ?Acute pancreatitis risk |

Based on findings from the DECLARE-TIMI 58 Trial, the benefit of SGLT-2 inhibitors for CHF/CKD is revised to read: "Benefit: empagliflozin, canagliflozin, dapagliflozin"

Choice of therapy after metformin

Drug-specific and patient factors to consider in adults with T2D

| | GLP-1RA Shorter-acting | GLP-1RA Long-acting | SGLT-2is | DPP-4is | Sulfonylureas | TZDs | Insulins |
|------------------------------------|--|--|--|--|---|--|---|
| Efficacy (↓HbA _{1c})* | Intermediate – High | High – very high | Intermediate–high (dependent on GFR) | Intermediate | High | High | Very high |
| Hypoglycaemia | No | No | No | Optimal strategy: <ul style="list-style-type: none"> regimens that stabilize hyperglycemia across multiple pathways act synergistically to reduce CV and other risk factors preserve β-cells | | | |
| Weight | Loss | Loss | Loss | | | | |
| CV effects | Improves cardiovascular risk factors | ↓ MACE with some agents | ↓ MACE, HF, CKD with some agents | | | | |
| Disadvantages/ adverse effects | Frequent GI side effects that may be transient | GI side effects, including gallbladder disease | Genital infections; risk dose adjustment/avoidance for renal disease | Rare urticaria/angioedema; increased risk of HF hospitalisation | Dose adjustment/avoidance for renal disease; high rate of secondary failure | Oedema/heart failure bone loss; bone fractures | Frequent dose adjustment for optimal efficacy |

*Davies MJ et al. *Diabetologia* 2018;61:2461–2498

THE JOURNAL OF CLINICAL AND APPLIED RESEARCH AND EDUCATION

VOLUME 43 | SUPPLEMENT 1

Diabetes Care.

WWW.DIABETES.ORG/DIABETESCARE

JANUARY 2018

SUPPLEMENT
1

AMERICAN DIABETES ASSOCIATION


STANDARDS OF MEDICAL CARE IN DIABETES—2020

 American
Diabetes
Association.
ISSN 0149-5992

ADA 2020 Consensus Report

COMPREHENSIVE MEDICAL EVALUATION AND ASSESSMENT OF COMORBIDITIES

DECISION CYCLE FOR PATIENT-CENTERED GLYCEMIC MANAGEMENT IN TYPE 2 DIABETES



ASCVD = Atherosclerotic Cardiovascular Disease

CKD = Chronic Kidney Disease

HF = Heart Failure

DSMES = Diabetes Self-Management Education and Support

SMBG = Self-Monitored Blood Glucose

GLYCEMIC TARGETS


Table 6.3—Summary of glycemic recommendations for many nonpregnant adults with diabetes

| | |
|---|--------------------------------|
| A1C | <7.0% (53 mmol/mol)* |
| Preprandial capillary plasma glucose | 80–130 mg/dL* (4.4–7.2 mmol/L) |
| Peak postprandial capillary plasma glucose† | <180 mg/dL* (10.0 mmol/L) |

*More or less stringent glycemic goals may be appropriate for individual patients. Goals should be individualized based on duration of diabetes, age/life expectancy, comorbid conditions, known CVD or advanced microvascular complications, hypoglycemia unawareness, and individual patient considerations. †Postprandial glucose may be targeted if A1C goals are not met despite reaching preprandial glucose goals. Postprandial glucose measurements should be made 1–2 h after the beginning of the meal, generally peak levels in patients with diabetes.

GLYCEMIC TARGETS

Approach to Individualization of Glycemic Targets





Diabetes and Sepsis: Risk, Recurrence, and Ruination

Lynn M. Frydrych¹, Fatemeh Fattahi², Katherine He¹, Peter A. Ward¹
and Matthew J. Delano^{1*}

**Διαβήτης και Σήψη: Κίνδυνος, Υποτροπή
και Καταστροφή**


Diabetes and Sepsis: Risk, Recurrence, and Ruination

- **A series of dysregulated physiological responses are generated, leading to organ dysfunction and a 10% mortality risk.**
- **Type II diabetes is a common and devastating disease** frequently encountered by clinicians who care for critically ill patients.
- Patients with **T2D have an increased risk of developing infections and sepsis**
- **T2D also worsens infection prognosis**, with T2D patients showing increased morbidity and mortality from sepsis

Diabetes and Sepsis: Risk, Recurrence, and Ruination


- Ερευνητές έχουν επικεντρώσει τις προσπάθειές τους στην κατανόηση των υποκείμενων διαταραχών του ανοσοποιητικού συστήματος που διευκολύνουν την ανάπτυξη επιπλοκών, δυσχεραίνουν την ανάρρωση από τη σήψη και αυξάνουν μακροχρόνια τη θνησιμότητα
- Ωστόσο, απαιτείται επικέντρωση και περισσότερη γνώση στην αλληλεπίδραση μεταξύ T2D, σήψης, ανοσίας και του αντίκτυπου τους στη συνολική επιβίωση.

Immune dysregulation in Type II diabetes and sepsis



- Ο διαβήτης χαρακτηρίζεται από ανοσολογική ανεπάρκεια, χρόνια φλεγμονή και ανοσοκαταστολή επηρεάζοντας έτσι τη συνολική ομοιόσταση του ανοσοποιητικού συστήματος.
- Επιπλέον, μελέτες έχουν δείξει ότι μια διαρκής φλεγμονώδης κατάσταση που οφείλεται σε δυσλειτουργία της ανοσίας, καταλήγει σε τραυματισμό οργάνων και θάνατο ασθενών.
- Όλα αυτά οδήγησαν σε υποθέσεις ότι τελικά αυτές οι επίμονες διαταραχές της λειτουργίας του ανοσοποιητικού συστήματος είναι οι κύριοι συντελεστές αυτής της αυξημένης θνησιμότητας.


Innate vs adaptive immune responses in sepsis and Type II diabetes



- Κατά τη διάρκεια ενός οξέος επεισοδίου σηψαιμίας, το ανοσοποιητικό σύστημα βρίσκεται σε μια συνεχή κατάσταση διακύμανσης
- Αυτό το συνεχές φαινόμενο της “τραμπάλας” θεωρείται ότι οδηγεί σε συνεχή φλεγμονή, διευκολύνει τον τραυματισμό των οργάνων και επιτρέπει την εμφάνιση επιπλοκών

Innate vs adaptive immune responses in sepsis and Type II diabetes

- Στον διαβήτη, το ανοσοποιητικό σύστημα βιώνει χρόνιες διαταραχές που οφείλονται στη χρόνια φλεγμονή, τοποθετώντας έτσι αυτά τα συστήματα σε μια συνεχή ροή
- Όταν αυτά τα δύο συστήματα (σήψη και διαβήτης) επικαλύπτονται, οι ασθενείς έχουν αυξημένη νοσηρότητα και θνησιμότητα
- Ωστόσο, υπάρχουν ασάφειες σχετικά με τις συνεργικές και ανταγωνιστικές αλλαγές που συμβαίνουν και τελικά οδηγούν σε επιδείνωση των διαταραχών του ανοσοποιητικού συστήματος και στην αδυναμία επιστροφής στην ομοιόσταση



Diabetes and Sepsis: Risk, Recurrence, and Ruination

- Ο διαβήτης τύπου II είναι μια ασθένεια με διαταραχές της ανοσίας που οδηγεί σε παρατεταμένη φλεγμονή, καταστολή του ανοσοποιητικού συστήματος και σημαντική νοσηρότητα
- Κλινικά, είναι προφανές ότι οι ασθενείς με T2D είναι πιο ευαίσθητοι σε λοιμώξεις
- Στη σήψη, παρά τις καλύτερες θεραπείες με στόχο τον έλεγχο της υπεργλυκαιμίας, τη χορήγηση αντιβιοτικών νωρίς και την πρόληψη της βλάβης των οργάνων, οι ασθενείς με T2D εξακολουθούν να έχουν χειρότερη νοσηρότητα και θνησιμότητα για λόγους που είναι ελάχιστα κατανοητοί
- Ωστόσο, η σχέση μεταξύ των δύο φαίνεται να είναι τα μη ρυθμιζόμενα ανοσοποιητικά μονοπάτια
- Τελικά οι ανοσορυθμιστικές θεραπείες που εισάγονται με πολύ στρατηγικό τρόπο και επηρεάζουν τις αλληλεπικαλυπτόμενες ανοσολογικές διαταραχές μεταξύ αυτών των δύο ασθενειών, έχουν τη δυνατότητα να βελτιώσουν ουσιαστικά τη συνολική νοσηρότητα και θνησιμότητα που αντιμετωπίζουν αυτά τα άτομα

Section 15.

Diabetes Care in the Hospital

DIABETES CARE IN THE HOSPITAL

Hospital Care Delivery Standards

- 15.1 Perform an A1C test on all patients with diabetes or hyperglycemia (blood glucose >140 mg/dL) admitted to the hospital if not performed in the prior 3 months. **B**
- 15.2 Insulin should be administered using validated written or computerized protocols that allow for predefined adjustments in the insulin dosage based on glycemic fluctuations. **C**

DIABETES CARE IN THE HOSPITAL

Diabetes Care Providers in the Hospital

- 15.3** When caring for hospitalized patients with diabetes, consult with a specialized diabetes or glucose management team when possible. **C**

DIABETES CARE IN THE HOSPITAL

Glycemic Targets in Hospitalized Patients

- 15.4** **Insulin therapy** should be initiated for **treatment of persistent hyperglycemia** starting at a threshold ≥ 180 mg/dL. Once insulin therapy is started, a target glucose range of **140–180 mg/dL is recommended** for the majority of critically ill patients and noncritically ill patients. **A**
- 15.5** More stringent goals, such as 110–140 mg/dL, may be appropriate for selected patients if they can be achieved without significant hypoglycemia. **C**

DIABETES CARE IN THE HOSPITAL

Hypoglycemia

- 15.8 A hypoglycemia management protocol should be adopted and implemented by each hospital or hospital system. A plan for preventing and treating hypoglycemia should be established for each patient. Episodes of hypoglycemia in the hospital should be documented in the medical record and tracked. **E**
- 15.9 The treatment regimen should be reviewed and changed as necessary to prevent further hypoglycemia when a blood glucose value of <70 mg/dL is documented. **C**

DIABETES CARE IN THE HOSPITAL

Glucose-lowering Treatment in Hospitalized Patients

15.6 Basal insulin or a basal plus bolus correction insulin regimen is the preferred treatment for noncritically ill hospitalized patients with poor oral intake or those who are taking nothing by mouth. **A**

An insulin regimen with basal, prandial, and correction components is the preferred treatment for non-critically ill hospitalized patients with good nutritional intake. **A**

15.7 Use of only a sliding scale insulin regimen in the inpatient hospital setting is strongly discouraged. **A**

Υπεργλυκαιμία & Βαρέως Πάσχων

- Η ρύθμιση του σακχάρου αίματος είναι απαραίτητη στους ασθενείς με σήψη
- ↑ επίπτωση θανάτου στους ασθενείς με υπεργλυκαιμία από stress σε σχέση με ασθενείς με φυσιολογικό σάκχαρο αίματος και σε σχέση με διαβητικούς
- ↑ επίπτωση στους διαβητικούς σε σχέση με τους μη διαβητικούς

Γλυκαιμικοί στόχοι στη ΜΕΘ..

- ΟΧΙ ΥΠΟΓΛΥΚΑΙΜΙΑ
- Έναρξη αντιμετώπισης αν $\text{Glu} > 180 \text{ mg/dl}$
- Επιθυμητό εύρος 140-180 mg/dl
- Σε υποπληθυσμούς (ΑΕΕ-ΚΕΚ-ΣΔ): όχι $\text{Glu} < 110 \text{ mg/dl}$


Σακχαρώδης Διαβήτης σε Νοσηλευόμενους Ασθενείς

- Οι ασθενείς με ΣΔ έχουν μεγαλύτερη πιθανότητα παρατεταμένης νοσηλείας συγκριτικά με μη-διαβητικούς ασθενείς
- Η υπεργλυκαιμία αλλά και η υπογλυκαιμία στη διάρκεια της νοσηλείας στο νοσοκομείο σχετίζεται με αυξημένη νοσηρότητα & θνητότητα



Τι Ορίζουμε ως Διαταραχές της Γλυκαιμίας σε Νοσηλευόμενους

- Εμμένουσα υπεργλυκαιμία > 140 mg/dl
 - HbA1c > 6.5 % υποδηλώνει ύπαρξη ΣΔ πριν τη νοσηλεία
 - Συγκέντρωση γλυκόζης < 70 mg/dl θεωρείται προειδοποιητική για πιθανή υπογλυκαιμία και απαιτεί τιτλοποίηση αγωγής
-
- Κλινικά σημαντική υπογλυκαιμία < 54 mg/dl
 - Κλινικά σοβαρή υπογλυκαιμία: σοβαρή γνωσιακή διαταραχή ανεξαρτήτως επιπέδων γλυκόζης



Δυσκολίες στη Διαχείριση του ΣΔ κατά τη Νοσηλεία

- Ώρες και ποσότητα γευμάτων
- Ώρες χορήγησης φαρμακευτικής αγωγής και συντονισμός χορήγησης με τα γεύματα
- Απουσία φυσικής δραστηριότητας
- Συνοδά νοσήματα και φαρμακευτικές αγωγές

Που Στοχεύει η Θεραπεία σε Νοσηλεύομενους Διαβητικούς Ασθενείς

ΣΤΟΧΕΥΕΙ

- Μείωση θνησιμότητας
- Μείωση ενδοноσοκομειακών λοιμώξεων
- Αποφυγή βαριάς υπεργλυκαιμίας και ΔΚΟ
- Μείωση μετεγχειρητικών επιπλοκών
- Φυσική επούλωση τραυμάτων, ελκών, κατακλίσεων
- Αποφυγή επεισοδίων υπογλυκαιμίας

Γλυκαιμικοί Στόχοι σε Νοσηλευόμενους Ασθενείς

Reviews/Commentaries/ADA Statements
CONSENSUS STATEMENT



American Association of Clinical Endocrinologists and American Diabetes Association Consensus Statement on Inpatient Glycemic Control

| Ασθενής | Στόχος σακχάρου αίματος (mg/dl) |
|-----------|--------------------------------------|
| Στην ΜΕΘ | 140 - 180 |
| Στο Τμήμα | Πριν το φαγητό: <140 Τυχαία: <180 |

Σε επιλεγμένους μη βαρέως πάσχοντες πιο αυστηροί στόχοι (110 - 140 mg/dl) χωρίς όμως υπογλυκαιμίες

Intensive versus Conventional Glucose Control in Critically Ill Patients

Τυχαιοποιημένη, προοπτική, μη τυφλή κλινική μελέτη με 6104 ασθενείς



- Καμμία διαφορά στη θνησιμότητα στις 30 ημέρες νοσηλείας
- Αύξηση της θνησιμότητας στις 90 ημέρες νοσηλείας στην ομάδα εντατικοποιημένης θεραπείας καθώς και περισσότερους θανάτους από καρδιαγγειακά συμβάματα και περισσότερες υπογλυκαιμίες

Intensive versus Conventional Glucose Control in Critically Ill Patients

Τυχαιοποιημένη, προοπτική, μη τυφλή κλινική μελέτη με 6104 ασθενείς

BOTTOM LINE



Glucose Goal: 140-180

- **Intensive** glycemic **control** (target 81-108 mg/dl) **increased deaths** compared to conventional control (targets ≤ 180 resulted in lower mortality than a target 81-108)

Online Submissions: wjg.wjgnet.com
wjg@wjgnet.com
doi:10.3748/wjg.15.4132



World J Gastroenterol 2009 September 7; 15(33): 4132-4136
World Journal of Gastroenterology ISSN 1007-9327
© 2009 The WJG Press and Baishideng. All rights reserved.

TOPIC HIGHLIGHT

Kazuhiro Hanazaki, MD, Professor and Chairman, *Series Editor*


Blood glucose control in patients with severe sepsis and septic shock

Hiroyuki Hirasawa, Shigeto Oda, Masataka Nakamura

Έλεγχος Γλυκόζης σε Ασθενείς με Σοβαρή Σήψη και Σηπτικό shock


- Οι εξελίξεις στη μοριακή βιολογία έχουν συμβάλει στην τεράστια πρόοδο της κατανόησης της παθοφυσιολογίας της σήψης.
- Τώρα είναι ευρέως αποδεκτό ότι τα κύρια χαρακτηριστικά της σήψης είναι η ανεξέλεγκτη ενεργοποίηση όχι μόνο των προ-φλεγμονωδών, αλλά και των αντιφλεγμονωδών αντιδράσεων, λόγω της συντριπτικής παραγωγής προ-φλεγμονωδών και αντιφλεγμονωδών μεσολαβητών. Αυτό προκαλεί πολλές παθολογικές αλλαγές σε ζωτικά όργανα και συστήματα, συμπεριλαμβανομένων και των μεταβολικών αλλαγών
- **Μία τέτοια μεταβολική αλλαγή είναι η υπεργλυκαιμία που προκύπτει από τη γλυκόλυση στους μυς και τη λιπόλυση, και την επακόλουθη γλυκονεογένεση και γλυκόλυση στο ήπαρ**

Illness leads to stress hyperglycemia




Stress hyperglycemia may also exacerbate illness.

The Sepsis Induced Changes in Circulating Hormones and Metabolites



Metabolic and Hormonal Changes Leading to Stress Hyperglycemia




Έλεγχος Γλυκόζης σε Ασθενείς με Σοβαρή Σήψη και Σηπτικό shock

- Υπάρχουν πολλές παθοφυσιολογικές διαταραχές κατά τη διάρκεια μιας σοβαρής σήψης ή/και σηπτικού shock, και μία από τις πιο εντυπωσιακές είναι η μεταβολική διαταραχή. Μεταξύ των μεταβολικών αλλαγών, η υπεργλυκαιμία είναι η πιο σημαντική
- Ο **Van den Berghe** και συν. έδειξαν τη σπουδαιότητα αντιμετώπισης της υπεργλυκαιμίας σε σοβαρή σήψη και στο σηπτικό shock και τα αποτελέσματα αυτής της μελέτης επισημαίνουν την αποτελεσματικότητα της εντατικής θεραπείας με ινσουλίνη στον γλυκαιμικό έλεγχο
- Ωστόσο, μια πιο **πρόσφατη** τυχαίοποιημένη **μελέτη** έδειξε ότι ένας τέτοιος γλυκαιμικός έλεγχος δεν είναι αποτελεσματικός στη μείωση της θνησιμότητας στην ICU και ότι ο γλυκαιμικός έλεγχος με εντατική θεραπεία με ινσουλίνη αυξάνει τον κίνδυνο υπογλυκαιμίας και τις επιπλοκές που προκύπτουν από την υπογλυκαιμία

Έλεγχος Γλυκόζης σε Ασθενείς με Σοβαρή Σήψη και Σηπτικό shock

- Η υπεργλυκαιμία σε σοβαρές παθήσεις, όπως η σήψη, δεν είναι μόνο ένας δείκτης σοβαρότητας της πάθησης και προγνωστικός παράγοντας κακής έκβασης, αλλά έχει και πολλές δυσμενείς επιπτώσεις σε ζωτικά όργανα.
- Μια τέτοια δυσμενή επίδραση στο ανοσοποιητικό σύστημα βλάπτει την ικανότητα του ξενιστή να καταπολεμήσει τη μόλυνση, με αποτέλεσμα τη μειωμένη δραστηριότητα των ουδετερόφιλων αλλά και μεταβολών στις κυτοκίνες, με αυξημένες συγκεντρώσεις των προφλεγμονωδών κυτοκινών TNF- α και IL -6, καθώς και μείωση παραγωγής του ενδοθηλιακού νιτρικού οξειδίου





Το ποσοστό επιτυχίας του εντατικού γλυκαιμικού ελέγχου ήταν σχετικά υψηλό στους ασθενείς με IL-6 <1000 pg / mL κατά την εισαγωγή στην ICU ενώ ήταν πολύ χαμηλό μεταξύ των ασθενών των οποίων το αρχικό επίπεδο IL-6 ήταν > 10.000 pg / mL

Έλεγχος Γλυκόζης σε Ασθενείς με Σοβαρή Σήψη και Σηπτικό shock


- Το 2001 ο Van den Berghe ανέφερε ότι σε ασθενείς ICU, η διατήρηση επιπέδων γλυκόζης μεταξύ 80 και 110 mg / dL με εντατική θεραπεία με ινσουλίνη, είχε ως αποτέλεσμα τη βελτίωση της επιβίωσης και της διάρκειας παραμονής τους στο νοσοκομείο
- Οι ίδιοι ερευνητές ανέφεραν αργότερα ότι η εντατική θεραπεία με ινσουλίνη μείωσε τη νοσηρότητα αλλά όχι τη θνησιμότητα στις ICU (Van den Berghe 2006)
- Αυτές οι μελέτες επηρέασαν και τις κατευθυντήριες οδηγίες για τη διαχείριση της σοβαρής σήψης και του σηπτικού shock ώστε να συστήσουν τον αυστηρό γλυκαιμικό έλεγχο ως μία από τις πιο σημαντικές θεραπευτικές προσεγγίσεις

More intensive glucose control with IV insulin resulted in lower ICU and in-hospital mortality

Intensive Insulin Therapy in the Surgical ICU: The Leuven Study




Hyperglycemia: An Independent Marker of In-Hospital Mortality in Patients with Undiagnosed Diabetes



.....all hospitalized patients should be screened for hyperglycemia

The immune response of patients with diabetes mellitus during sepsis




During early sepsis patients with T1D have:

- defective neutrophil functions, resulting in impaired bacterial control at the site of infection, bacteremia and a poor sepsis outcome

The immune response of patients with diabetes mellitus during sepsis

Patients with T2D seem to have a better sepsis prognosis due to several reasons:

- diets rich in unsaturated fat
- use of prebiotics
- higher levels of leptin and
- the use of PPAR γ agonists



AACE-ADA Consensus Statement on Inpatient Glycaemic Targets

Recommends using insulin therapy if blood glucose levels exceed 180 mg/dL

Non-ICU Setting (general medical/surgical)

- Subcutaneous insulin preferred
- Target BG levels:
 - Pre-meal BG <140 mg/dL
 - Random BG <180 mg/dL
- More stringent targets in stable patients
- Less stringent targets in patients with severe comorbidities

ICU Setting

- Insulin infusion preferred
- Starting threshold not higher than 180 mg/dL
- Maintain between 140–180 mg/dL
- Blood glucose <110 mg/dL is not recommended

AACE/ADA Consensus Statement

Inpatient glycemic targets

- Insulin infusion to control hyperglycemia
- Starting threshold no higher than 180 mg/dl
- Maintain BG between 140 and 180 mg/dl
 - Possible greater benefit at lower end of range
- Somewhat lower targets may be appropriate in selected patients
- Targets <110 mg/dl are not recommended

| | | | |
|---|---|--|---|
| Not recommended < 110 mg/dl | May be appropriate 110-140 mg/dl | Recommended 140-180 mg/dl | Not recommended > 180 mg/dl |
|---|---|--|---|

New AACE-ADA Consensus Statement on Inpatient Glycemic Control

ICU Setting

- Insulin infusion preferred
- Starting threshold not higher than 180 mg/dl
- Maintain BG **140–180 mg/dl**
(greater benefit likely at *lower end of this range*)
- Lower targets (not evidence-based) may be appropriate in selected patients if already being successfully achieved
- <110 NOT recommended (not safe)

Non-ICU Setting

- Most patients:
 - **pre-meal BG <140 mg/dL**
 - **random BG <180 mg/dL**
- More stringent targets may be appropriate in stable patients
- Less stringent targets may be appropriate in patients with severe comorbidities
- Scheduled SQ insulin with basal-nutritional-correction preferred; avoid prolonged therapy RISS alone

AACE, American Association of Clinical Endocrinologists; ADA, American Diabetes Association; SQ, subcutaneous; RISS, regular insulin sliding scale.

CONCLUSION

It is now suggested

- ...**tight glycemic control** with a target BG level of 90-110 mg/dL does **not improve clinical outcome**
- ..**less strict glycemic control** with a target BG level of 140-180 mg/dL is **more effective**
- Also **specific targeting** of glycemic control in diabetic patients should be considered
- ...**avoid hypoglycemia** during insulin therapy
- ..**control of hypercytokinemia** should be considered for more effective glycemic control in patients with severe sepsis and septic shock



Thank you for your attention!



QUESTIONS – COMMENTS

Discussion